

High-Dimensional Matched Subspace Detection When Data are Missing

Laura Balzano, Benjamin Recht, and Robert Nowak
University of Wisconsin-Madison

Abstract—We consider the problem of deciding whether a highly incomplete signal lies within a given subspace. This problem, Matched Subspace Detection, is a classical, well-studied problem when the signal is completely observed. High-dimensional testing problems in which it may be prohibitive or impossible to obtain a complete observation motivate this work. The signal is represented as a vector in \mathbb{R}^n , but we only observe $m \ll n$ of its elements. We show that reliable detection is possible, under mild incoherence conditions, as long as m is slightly greater than the dimension of the subspace in question.

I. INTRODUCTION

Testing whether a signal lies within a subspace is a problem arising in a wide range of applications including medical [1] and hyperspectral [5] imaging, communications [6], radar [8], and anomaly detection [12]. The classical formulation of this problem is a binary hypothesis test of the following form. Let $v \in \mathbb{R}^n$ denote a signal and let $x = v + w$, where w is a noise of known distribution. We are given a subspace $S \subset \mathbb{R}^n$ and we wish to decide if $v \in S$ or not, based on x . Tests are usually based on some measure of the energy of x in the subspace S , and these ‘matched subspace detectors’ enjoy optimal properties [10], [11].

This paper considers a variation on this classical problem, motivated by high-dimensional applications where it is prohibitive or impossible to measure v completely. We assume that only a small subset $\Omega \subset \{1, \dots, n\}$ of the elements of v are observed (with or without noise), and based on these observations we want to test whether $v \in S$. For example, consider monitoring a large networked system such as a portion of the Internet. Measurement nodes in the network may have software that collects measurements such as upload and download rate, number of packets, or type of traffic given by the packet headers. In order to monitor the network these measurements will be collected in a central place for compilation, modeling and analysis. The effective dimension of the state of such systems is often much lower than the extrinsic dimension of the network itself. Subspace detection, therefore, can be a useful tool for detecting changes or anomalies. The challenge is that it may be impossible to obtain every measurement from every point in the network due to resource constraints, node outages, etc.

The main result of this paper answers the following question. Given a subspace S of dimension $r \ll n$, how many elements of v must be observed so that we can reliably decide if it belongs to S ? The answer is that, under some mild incoherence conditions, the number is just slightly larger than

r . This means that reliable matched subspace detectors can be constructed from very few measurements, making them scalable and applicable to large-scale testing problems.

The main focus of this paper is an estimator of the energy of v in S based on only observing the elements $\{v_i\}_{i \in \Omega}$. Section II proposes the estimator. Section III presents a theorem giving quantitative bounds on the estimator’s performance and the proof using three lemmas that are proved in the Appendix. Section IV presents numerical experiments. Section V applies the main result to the subspace detection problem, both with and without noise.

II. ENERGY ESTIMATION FROM INCOMPLETE DATA

Let v_Ω be the vector of dimension $|\Omega| \times 1$ comprised of the elements v_i , $i \in \Omega$, ordered lexicographically; here $|\Omega|$ denotes the cardinality of Ω . The energy of v in the subspace S is $\|P_S v\|_2^2$, where P_S denotes the projection operator onto S . There are two natural estimators of $\|P_S v\|_2^2$ based on v_Ω . The first is simply to form the $n \times 1$ vector \tilde{v} with elements v_i if $i \in \Omega$ and zero if $i \notin \Omega$, for $i = 1, \dots, n$. This ‘zero-filled’ vector yields the simple estimator $\|P_S \tilde{v}\|_2^2$. Filling missing elements with zero is a fairly common, albeit naïve, approach to dealing with missing data. Unfortunately, the estimator $\|P_S \tilde{v}\|_2^2$ is fundamentally flawed. Even if $v \in S$, the zero-filled vector \tilde{v} does not necessarily lie in S .

A better estimator can be constructed as follows. Let U be an $n \times r$ matrix whose columns span the r -dimensional subspace S . Note that for any such U , $P_S = U(U^T U)^{-1} U^T$. With this representation in mind, let U_Ω denote the $|\Omega| \times r$ matrix, whose rows are the $|\Omega|$ rows of U indexed by the set Ω , arranged in lexicographic order. Since we only observe v on the set Ω , another approach to estimating its energy in S is to assess how well v_Ω can be represented in terms of the rows of U_Ω . Define the projection operator $P_{S_\Omega} := U_\Omega(U_\Omega^T U_\Omega)^{-1} U_\Omega^T$. It follows immediately that if $v \in S$, then $\|v - P_S v\|_2^2 = 0$ and $\|v_\Omega - P_{S_\Omega} v_\Omega\|_2^2 = 0$, whereas $\|\tilde{v} - P_S \tilde{v}\|_2^2$ can be significantly greater than zero. This property makes $\|P_{S_\Omega} v_\Omega\|_2^2$ a much better candidate estimator than $\|P_S \tilde{v}\|_2^2$. However, if $|\Omega| \leq r$, then it is possible that $\|v_\Omega - P_{S_\Omega} v_\Omega\|_2^2 = 0$, even if $\|v - P_S v\|_2^2 > 0$. Our main result shows that if $|\Omega|$ is just slightly greater than r , then with high probability $\|v_\Omega - P_{S_\Omega} v_\Omega\|_2^2$ is very close to $\frac{|\Omega|}{n} \|v - P_S v\|_2^2$.

III. MAIN THEOREM

Let us now focus on our main goal of detecting from a very small number of samples whether there is energy in a vector

v outside the r -dimensional subspace S . In order to do so, we must first quantify how much information we can expect each sample to provide. The authors in [3] defined the *coherence* of a subspace S to be the quantity

$$\mu(S) := \frac{n}{r} \max_j \|P_S e_j\|_2^2.$$

That is, $\mu(S)$ measures the maximum magnitude attainable by projecting a standard basis element onto S . Note that $1 \leq \mu(S) \leq \frac{n}{r}$. The minimum $\mu(S) = 1$ can be attained by looking at the span of any r columns of the discrete Fourier transform. Any subspace that contains a standard basis element will maximize $\mu(S)$. For a vector z , we let $\mu(z)$ denote the coherence of the subspace spanned by z . By plugging in the definition, we have

$$\mu(z) = \frac{n\|z\|_\infty^2}{\|z\|_2^2}.$$

To state our main theorem, write $v = x + y$ where $x \in S$ and $y \in S^\perp$. Let the entries of v be sampled uniformly with replacement, and again let Ω refer to the set of indices for observations of entries in v . Given these conventions, we have the following.

Theorem 1. *Let $\delta > 0$ and $m \geq \frac{8}{3}r\mu(S) \log(\frac{2r}{\delta})$. Then with probability at least $1 - 4\delta$,*

$$\frac{m(1-\alpha) - r\mu(S) \frac{(1+\beta)^2}{(1-\gamma)}}{n} \|v - P_S v\|_2^2 \leq \|v_\Omega - P_{S_\Omega} v_\Omega\|_2^2$$

and

$$\|v_\Omega - P_{S_\Omega} v_\Omega\|_2^2 \leq (1+\alpha) \frac{m}{n} \|v - P_S v\|_2^2$$

where $\alpha = \sqrt{\frac{2\mu(y)^2}{m} \log(\frac{1}{\delta})}$, $\beta = \sqrt{2\mu(y) \log(\frac{1}{\delta})}$, and $\gamma = \sqrt{\frac{8r\mu(S)}{3m} \log(\frac{2r}{\delta})}$.

Proof: In order to prove the theorem, we split the quantity of interest into three terms and bound each with high probability. Consider $\|v_\Omega - P_{S_\Omega} v_\Omega\|_2^2 = \|y_\Omega - P_{S_\Omega} y_\Omega\|_2^2$. Let the r columns of U be an orthonormal basis for the subspace S . We want to show that

$$\|y_\Omega - P_{S_\Omega} y_\Omega\|_2^2 = \|y_\Omega\|_2^2 - y_\Omega^T U_\Omega (U_\Omega^T U_\Omega)^{-1} U_\Omega^T y_\Omega \quad (1)$$

is near $\frac{m}{n} \|y\|_2^2$ with high probability. To proceed, we need the following three Lemmas whose proofs can be found in the Appendix.

Lemma 1. *With the same notations as Theorem 1,*

$$(1-\alpha) \frac{m}{n} \|y\|_2^2 \leq \|y_\Omega\|_2^2 \leq (1+\alpha) \frac{m}{n} \|y\|_2^2$$

with probability at least $1 - 2\delta$.

Lemma 2. *With the same notations as Theorem 1,*

$$\|U_\Omega^T y_\Omega\|_2^2 \leq (\beta+1)^2 \frac{m}{n} \frac{r\mu(S)}{n} \|y\|_2^2$$

with probability at least $1 - \delta$.

Lemma 3. *With the same notations as Theorem 1,*

$$\|(U_\Omega^T U_\Omega)^{-1}\|_2 \leq \frac{n}{(1-\gamma)m}$$

with probability at least $1 - \delta$, provided that $\gamma < 1$.

To apply these three Lemmas, write the second term of Equation (1) as

$$y_\Omega^T U_\Omega (U_\Omega^T U_\Omega)^{-1} U_\Omega^T y_\Omega = \|W_\Omega U_\Omega^T y_\Omega\|_2^2$$

where $W_\Omega^T W_\Omega = (U_\Omega^T U_\Omega)^{-1}$. By Lemma 3, $U_\Omega^T U_\Omega$ is invertible under the assumptions of our theorem, and hence W_Ω is well-defined and has spectral norm bounded by the square root of the inverse of the smallest eigenvalue of $U_\Omega^T U_\Omega$. That is, we have

$$\begin{aligned} \|W_\Omega U_\Omega^T y_\Omega\|_2^2 &\leq \|W_\Omega\|_2^2 \|U_\Omega^T y_\Omega\|_2^2 \\ &= \|W_\Omega^T W_\Omega\|_2 \|U_\Omega^T y_\Omega\|_2^2 \\ &= \|(U_\Omega^T U_\Omega)^{-1}\|_2 \|U_\Omega^T y_\Omega\|_2^2. \end{aligned}$$

$\|(U_\Omega^T U_\Omega)^{-1}\|_2$ is bounded by Lemma 3 and $\|U_\Omega^T y_\Omega\|_2$ is bounded by Lemma 2. Putting these two bounds together with the bounds in Lemma 1 and using the union bound, we have that with probability at least $1 - 4\delta$

$$\begin{aligned} (1+\alpha)^2 \frac{m}{n} \|y\|_2^2 &\geq \|y_\Omega\|_2^2 - \|(U_\Omega^T U_\Omega)^{-1}\|_2 \|U_\Omega^T y_\Omega\|_2^2 \\ &\geq (1-\alpha)^2 \frac{m}{n} \|y\|_2^2 - \frac{(\beta+1)^2 r\mu(S)}{(1-\gamma)n} \|y\|_2^2 \end{aligned}$$

giving us our bound. \blacksquare


IV. DISCUSSION AND NUMERICAL EXPERIMENTS

In this section we wish to give some intuition for the lower bound in Theorem 1 and show what the estimate $\|v_\Omega - P_{S_\Omega} v_\Omega\|_2$ looks like in simulations. If we are able to make the parameters α, β, γ small, we have a lower bound which looks like

$$\frac{m - r\mu(S)}{n} \|v - P_S v\|_2$$

For an incoherent subspace, the parameter $\mu(S) = 1$. In this case the result says that for $m \leq r$ the bound is ≤ 0 , which is consistent with the fact that $\|v_\Omega - P_{S_\Omega} v_\Omega\|_2 = 0$ always for $m \leq r$. However once $m \geq r + 1$, linear algebraic reasoning tells us that $\|v_\Omega - P_{S_\Omega} v_\Omega\|_2$ will be strictly positive with positive probability; Theorem 1 goes further to say the norm is strictly positive with high probability once $m \sim \mathcal{O}(r \log r)$.

The parameters α, β, γ all depend on $\sqrt{\log(\frac{1}{\delta})}$, so as we crank down on δ we must allow for very low probability situations where the norm $\|v_\Omega - P_{S_\Omega} v_\Omega\|_2$ is not close to $\frac{m}{n} \|v - P_S v\|_2$. For α and γ we can compensate by increasing m , but that doesn't help β . In fact, even if the vector y is incoherent and $\mu(y) = 1$, its minimum value, then $\beta = 2$ for $\delta \approx .135$. To get β very near zero, δ must be very near one, but this is not a useful regime. It turns out that in expectation [2], the parameters α, β play no role, and we have that



(a) Incoherent subspace (spanned by the columns of the Fourier basis). $r = 10$, $n = 100$, $\mu(S) = 1.815$, $\mu(y) = 3.458$.
(b) Coherent subspace. $r = 10$, $n = 120$, $\mu(S) = 2.096$, $\mu(y) = 12.220$.

Fig. 1: Each of the simulations had a fixed vector and a particular sample size m but different sample set Ω , and we show the mean, minimum and maximum projection error $\|v_\Omega - P_{S_\Omega} v_\Omega\|_2^2$ of those 100 simulations.

$$\mathbb{E} [\|v_\Omega - P_{S_\Omega} v_\Omega\|_2^2] \geq \frac{m - \frac{r\mu(S)}{1-\gamma}}{n} \|v - P_S v\|_2^2$$

This plays out in simulations of Figure 1, where we see qualitatively that for very incoherent subspaces, $\|v_\Omega - P_{S_\Omega} v_\Omega\|_2$ is positive for $m > r\mu(S) \log r$. Our simulations for the Fourier basis and a basis made of orthogonalized Gaussian random vectors always showed the estimate to be positive for $m > r\mu(S) \log r$, even for the worst-case simulation run. For more coherent subspaces, we often (but not always) see that the norm is positive as long as $m > r\mu(S) \log r$.

We also want to note that the bound in Lemma 2 is very loose as m gets close to n . For m very small, the norm $\|U_\Omega^T y_\Omega\|_2$ will also be very small simply because very few entries are observed. As m grows, the norm will grow, but we know that for $m = n$ the norm $\|U_\Omega^T y_\Omega\|_2 = \|U^T y\|_2 = 0$ with probability 1, because y_Ω is orthogonal to the subspace. This nonlinear behavior as $m \rightarrow n$ makes for a loose lower bound in that regime.

V. MATCHED SUBSPACE DETECTION

We have the following detection set up. Our hypotheses are $\mathcal{H}_0 : v \in S$ and $\mathcal{H}_1 : v \notin S$ and the test statistic we will use is

$$t(v_\Omega) = \|v_\Omega - P_{S_\Omega} v_\Omega\|_2^2 \underset{\mathcal{H}_0}{\overset{\mathcal{H}_1}{\gtrless}} \eta$$

In the noiseless case, we can let $\eta = 0$; our result in Theorem 1 shows for $\delta > 0$, the probability of detection is $P_D = \mathbb{P}[t(v_\Omega) > 0 | \mathcal{H}_1] \geq 1 - 4\delta$ as long as m is large enough, and we also have that the probability of false alarm is zero, $P_{FA} = \mathbb{P}[t(v_\Omega) > 0 | \mathcal{H}_0] = 0$ since the projection error will be zero when $v \in S$.

When we introduce noise we have the same hypotheses, but we compute the statistic on $\tilde{v}_\Omega = v_\Omega + w$ where $w \sim \mathcal{N}(0, 1)$ is Gaussian white noise:

$$t(\tilde{v}_\Omega) = \|\tilde{v}_\Omega - P_{S_\Omega} \tilde{v}_\Omega\|_2^2 \underset{\mathcal{H}_0}{\overset{\mathcal{H}_1}{\gtrless}} \eta_\lambda$$




Fig. 2: Simulation results for the zero-filling approach, $v \in S$, $\|v\|_2^2 = 1$. The basis used is the Fourier basis, $r = 10$, $n = 120$, $\mu(S) = 1.771$, $\mu(y) = 3.808$. Note that the zero-filled error curves can be made arbitrarily large by increasing $\|v\|_2^2$.

We choose η_λ to fix the probability of false alarm:

$$\mathbb{P}[t(\tilde{v}_\Omega) > \eta_\lambda | \mathcal{H}_0] \leq \lambda = P_{FA}$$

And then we have from [10] that $t(\tilde{v}_\Omega)$ is distributed as a non-central χ^2 with r degrees of freedom and non-centrality parameter $\|v_\Omega - P_{S_\Omega} v_\Omega\|_2^2$, and that P_D is monotonically increasing with the non-centrality parameter. Putting this together with Theorem 1 we see that as m grows, $\|v_\Omega - P_{S_\Omega} v_\Omega\|_2^2$ grows and thus the probability of detection grows.

We now show why the heuristic approach of zero-filling the incomplete vector v_Ω does not work. As we described in Section II, the zero-filling approach is to fill the vector v with zeros and then project onto the full subspace S . We denote the zero-filled vector as v_0 and then calculate the projection energy only on the observed entries:

$$t_0(v_\Omega) = \|v_\Omega - (P_S v_0)_\Omega\|_2^2 \underset{\mathcal{H}_0}{\overset{\mathcal{H}_1}{\gtrless}} \eta$$

Simple algebraic consideration reveals that $t_0(v_\Omega) | \mathcal{H}_0$ is positive. In fact, even in the absence of noise, the probability of false alarm can be arbitrarily large as $\|v\|_2^2$ increases. The value of $t_0(v_\Omega) | \mathcal{H}_0$, based on noiseless observations, is plotted as a function of the number of measurements in Figure 2.

We note that for unknown noise power or structured interference, these results can be extended using the GLRT [11].

VI. CONCLUSION

We have shown that it is possible to detect whether a highly incomplete vector has energy outside a subspace. This is a fundamental result to add to a burgeoning collection of results for incomplete data analysis given a low-rank assumption. Missing data are the norm and not the exception in any massive data collection system, so this result has implications on many other areas of study. In future work it will be important to understand how crucial the sampling assumption is to the result. It would be interesting to address sampling without replacement and other sampling models. Additionally, understanding how the bound should look as m nears n will be interesting for applications where the vector is not too incomplete and the energy outside the subspace needs to be accurately estimated.

APPENDIX

We will need the following two large deviation bounds in the proofs of our Lemmas below.

Theorem 2 (McDiarmid's Inequality [7]). *Let X_1, \dots, X_n be independent random variables, and assume f is a function for which there exist t_i , $i = 1, \dots, n$ satisfying*

$$\sup_{x_1, \dots, x_n, \hat{x}_i} |f(x_1, \dots, x_n) - f(x_1, \dots, \hat{x}_i, \dots, x_n)| \leq t_i$$

where \hat{x}_i indicates replacing the sample value x_i with any other of its possible values. Call $f(X_1, \dots, X_n) := Y$. Then for any $\epsilon > 0$,

$$\mathbb{P}[Y \geq \mathbb{E}[Y] + \epsilon] \leq \exp\left(\frac{-2\epsilon^2}{\sum_{i=1}^n t_i^2}\right) \quad (2)$$

$$\mathbb{P}[Y \leq \mathbb{E}[Y] - \epsilon] \leq \exp\left(\frac{-2\epsilon^2}{\sum_{i=1}^n t_i^2}\right) \quad (3)$$

Theorem 3 (Noncommutative Bernstein Inequality [4], [9]). *Let X_1, \dots, X_m be independent zero-mean square $r \times r$ random matrices. Suppose $\rho_k^2 = \max\{\|\mathbb{E}[X_k X_k^T]\|_2, \|\mathbb{E}[X_k^T X_k]\|_2\}$ and $\|X_k\|_2 \leq M$ almost surely for all k . Then for any $\tau > 0$,*

$$\mathbb{P}\left[\left\|\sum_{k=1}^m X_k\right\|_2 > \tau\right] \leq 2r \exp\left(\frac{-\tau^2/2}{\sum_{k=1}^m \rho_k^2 + M\tau/3}\right)$$

We now proceed with the proof of our three central Lemmas.

Proof of Lemma 1: To prove this we use McDiarmid's inequality from Theorem 2 for the function $f(X_1, \dots, X_m) = \sum_{i=1}^m X_i$. The resulting inequality is more commonly referred to as Hoeffding's inequality.

We begin with the first inequality. Set $X_i = y_{\Omega(i)}^2$. We seek a good value for t_i . Since $y_{\Omega(i)}^2 \leq \|y\|_\infty^2$ for all i , we have

$$\begin{aligned} |f(X_1, \dots, X_m) - f(X_1, \dots, \hat{X}_k, \dots, X_m)| &= |X_k - \hat{X}_k| \\ &\leq 2\|y\|_\infty^2 \end{aligned}$$

We calculate $\mathbb{E}[\sum_{i=1}^m X_i]$ as follows. Define \mathbb{I}_{Ω} to be the indicator function, and assume that the samples are taken uniformly with replacement.

$$\begin{aligned} \mathbb{E}\left[\sum_{i=1}^m X_i\right] &= \mathbb{E}\left[\sum_{i=1}^m y_{\Omega(i)}^2\right] \\ &= \sum_{i=1}^m \mathbb{E}\left[\sum_{j=1}^n y_j^2 \mathbb{I}_{\{\Omega(i)=j\}}\right] = \frac{m}{n} \|y\|_2^2 \end{aligned}$$

Plugging into Equation (3), the left hand side is

$$\mathbb{P}\left[\sum_{i=1}^m X_i \leq \mathbb{E}\left[\sum_{i=1}^m X_i\right] - \epsilon\right] = \mathbb{P}\left[\sum_{i=1}^m X_i \leq \frac{m}{n} \|y\|_2^2 - \epsilon\right]$$

and letting $\epsilon = \alpha \frac{m}{n} \|y\|_2^2$, we then have that this probability is bounded by

$$\exp\left(\frac{-2\alpha^2 \left(\frac{m}{n}\right)^2 \|y\|_2^4}{4m \|y\|_\infty^4}\right)$$

Thus, the resulting probability bound is

$$\mathbb{P}\left[\|y_{\Omega}\|_2^2 \geq (1 - \alpha) \frac{m}{n} \|y\|_2^2\right] \geq 1 - \exp\left(\frac{-\alpha^2 m \|y\|_2^4}{2n^2 \|y\|_\infty^4}\right) \quad (4)$$

Substituting our definitions of $\mu(y)$ and α shows that the lower bound holds with probability at least $1 - \delta$. The argument for the upper bound is identical after replacing Equation (2) instead of (3). The Lemma now follows by applying the union bound. ■

Proof of Lemma 2: We use McDiarmid's inequality in a very similar fashion to the proof of Lemma 1. Let $X_i = y_{\Omega(i)} U_{\Omega(i)}$, where $\Omega(i)$ refers to the i^{th} sample index. Thus $y_{\Omega(i)}$ is a scalar, and the notation $U_{\Omega(i)}$ refers to an $r \times 1$ vector representing the transpose of the $\Omega(i)^{\text{th}}$ row of U .

Let our function $f(X_1, \dots, X_m) = \|\sum_{i=1}^m X_i\|_2 = \|U_{\Omega}^T y_{\Omega}\|_2$. To find the t_i of the theorem we first need to bound $\|X_i\|$ for all i . Observe that $\|U_{\Omega(i)}\|_2 = \|U^T e_i\|_2 = \|P_S e_i\|_2 \leq \sqrt{r\mu(S)/n}$ by assumption. Thus,

$$\|X_i\|_2 \leq |y_{\Omega(i)}| \|U_{\Omega(i)}\|_2 \leq \|y\|_\infty \sqrt{r\mu(S)/n}$$

Then observe $|f(X_1, \dots, X_m) - f(X_1, \dots, \hat{X}_k, \dots, X_m)|$ is

$$\begin{aligned} \left\|\sum_{i=1}^m X_i\right\|_2 - \left\|\sum_{i \neq k} X_i + \hat{X}_k\right\|_2 &\leq \|X_k - \hat{X}_k\|_2 \\ &\leq \|X_k\|_2 + \|\hat{X}_k\|_2 \\ &\leq 2\|y\|_\infty \sqrt{\frac{r\mu(S)}{n}}. \end{aligned}$$

Here, the first two inequalities follow from the triangle inequality. Next we calculate a bound for $\mathbb{E}[f(X_1, \dots, X_m)] = \mathbb{E}[\|\sum_{i=1}^m X_i\|_2]$. Assume again that the samples are taken uniformly with replacement. We have

$$\sum_{k=1}^r U_{jk}^2 = \|P_S e_j\|^2 \leq \frac{r}{n} \mu(S),$$

from which we can see that

$$\begin{aligned} \mathbb{E}\left[\left\|\sum_{i=1}^m X_i\right\|_2^2\right] &= \mathbb{E}\left[\|U_{\Omega}^T y_{\Omega}\|_2^2\right] \\ &= \sum_{k=1}^r \mathbb{E}\left[\sum_{i=1}^m \sum_{j=1}^n U_{jk}^2 y_j^2 \mathbb{I}_{\{\Omega(i)=j\}}\right] \quad (5) \end{aligned}$$

$$\begin{aligned} &= \sum_{k=1}^r m \left(\sum_{j=1}^n U_{jk}^2 y_j^2\right) \frac{1}{n} \quad (6) \\ &\leq \frac{m}{n} \frac{r\mu(S)}{n} \|y\|_2^2 \end{aligned}$$

The step (5) follows because the cross terms cancel by orthogonality. The step (6) is because of our assumption that sampling is uniform with replacement.

Since $\mathbb{E}[\|X\|_2] \leq \mathbb{E}[\|X\|_2^2]^{1/2}$ by Jensen's inequality, we have that $\mathbb{E}[\|\sum_{i=1}^m X_i\|_2] \leq \sqrt{\frac{m}{n}} \sqrt{\frac{r\mu(S)}{n}} \|y\|_2$. Letting $\epsilon = \beta \sqrt{\frac{m}{n}} \sqrt{\frac{r\mu(S)}{n}} \|y\|_2$ and plugging into Equation (2), we then have that the probability is bounded by

$$\exp\left(\frac{-2\beta^2 \frac{m}{n} \frac{r\mu(S)}{n} \|y\|_2^2}{4m \|y\|_\infty^2 \frac{r\mu(S)}{n}}\right)$$

Thus, the resulting probability bound is

$$\mathbb{P}\left[\|U_\Omega y_\Omega\|_2^2 \leq (1 + \beta)^2 \frac{mr\mu(S)}{n^2} \|y\|_2^2\right] \geq 1 - \exp\left(\frac{-\beta^2 \|y\|_2^2}{2n \|y\|_\infty^2}\right)$$

Substituting our definitions of $\mu(y)$ and β shows that the lower bound holds with probability at least $1 - \delta$, completing the proof. \blacksquare

Proof of Lemma 3: We use the Noncommutative Bernstein Inequality as follows. Let $X_k = U_{\Omega(k)} U_{\Omega(k)}^T - \frac{1}{n} I_r$, where the notation $U_{\Omega(k)}$ is as before, i.e. is the transpose of the $\Omega(k)^{th}$ row of U , and I_r is the $r \times r$ identity matrix. Note that this random variable is zero mean.

We must compute ρ_k^2 and M . Since $\Omega(k)$ is chosen uniformly with replacement, the X_k are identically distributed, and ρ does not depend on k . For ease of notation we will denote $U_{\Omega(k)}$ as U_k .

Using the fact that for positive semi-definite matrices, $\|A - B\|_2 \leq \max\{\|A\|_2, \|B\|_2\}$, and recalling again that $\|U_k\|_2^2 = \|U^T e_k\|_2^2 = \|P_S e_k\|_2^2 \leq r\mu(S)/n$, we have

$$\left\|U_k U_k^T - \frac{1}{n} I_r\right\|_2 \leq \max\left\{\frac{r\mu(S)}{n}, \frac{1}{n}\right\}$$

and we let $M := r\mu(S)/n$.

For ρ , we note

$$\begin{aligned} \|\mathbb{E}[X_k X_k^T]\|_2 &= \|\mathbb{E}[X_k^T X_k]\|_2 \\ &= \left\|\mathbb{E}\left[\left(U_k U_k^T - \frac{1}{n} I_r\right)^2\right]\right\|_2 \\ &= \left\|\mathbb{E}\left[U_k U_k^T U_k U_k^T - \frac{2}{n} U_k U_k^T + \frac{1}{n^2} I_r\right]\right\|_2 \\ &= \left\|\mathbb{E}[U_k U_k^T U_k U_k^T] - \frac{1}{n^2} I_r\right\|_2 \\ &\leq \max\left\{\|\mathbb{E}[U_k U_k^T U_k U_k^T]\|, \frac{1}{n^2}\right\} \\ &\leq \max\left\{\frac{r\mu(S)}{n} \mathbb{E}\|U_k U_k^T\|_2, \frac{1}{n^2}\right\} \\ &= \max\left\{\frac{r\mu(S)}{n^2} \|I_r\|_2, \frac{1}{n^2}\right\} \\ &= \frac{r\mu(S)}{n^2}. \end{aligned}$$

Thus we let $\rho^2 := r\mu(S)/n^2$.

Now we can apply the Noncommutative Bernstein Inequality, Theorem 3. First we restrict τ to be such that $M\tau \leq m\rho^2$ to simplify the denominator of the exponent. Then we get that

$$2r \exp\left(\frac{-\tau^2/2}{m\rho^2 + M\tau/3}\right) \leq 2r \exp\left(\frac{-\tau^2/2}{\frac{4}{3}m \frac{r\mu(S)}{n^2}}\right)$$

and thus

$$\mathbb{P}\left[\left\|\sum_{k \in \Omega} \left(U_k U_k^T - \frac{1}{n} I_r\right)\right\| > \tau\right] \leq 2r \exp\left(\frac{-3n^2\tau^2}{8mr\mu(S)}\right)$$

Now take $\tau = \gamma m/n$ with γ defined in the statement of Theorem 1. Since $\gamma < 1$ by assumption, $M\tau \leq m\rho^2$ holds and we have

$$\mathbb{P}\left[\left\|\sum_{k \in \Omega} \left(U_k U_k^T - \frac{1}{n} I_r\right)\right\|_2 \leq \frac{m}{n} \gamma\right] \geq 1 - \delta$$

We note that $\left\|\sum_{k \in \Omega} U_k U_k^T - \frac{m}{n} I_r\right\|_2 \leq \frac{m}{n} \gamma$ implies that the minimum singular value of $\sum_{k \in \Omega} U_k U_k^T$ is at least $(1 - \gamma) \frac{m}{n}$. This in turn implies that

$$\left\|\left(\sum_{k \in \Omega} U_k U_k^T\right)^{-1}\right\|_2 \leq \frac{n}{(1 - \gamma)m}$$

which completes the proof. \blacksquare

REFERENCES

- [1] B. Ardekani, J. Kershaw, K. Kashikura, and I. Kanno. Activation detection in functional mri using subspace modeling and maximum likelihood estimation. *IEEE Transactions on Medical Imaging*, 18(2), February 1999.
- [2] L. Balzano, B. Recht, and R. Nowak. The energy of the projection of a vector with missing data is accurate in expectation. Technical report, Information Sciences Laboratory at the University of Wisconsin-Madison, November 2009.
- [3] E. Candès and B. Recht. Exact matrix completion via convex optimization. *Foundations of Computational Mathematics*, 9(6):717–772, December 2009.
- [4] D. Gross, Y.-K. Liu, S. T. Flammia, and J. E. S. Becker. Quantum state tomography via compressed sensing. 2009. Preprint available at <http://arxiv.org/abs/0909.3304v2>.
- [5] H. Kwon and N. Nasrabadi. Kernel matched subspace detectors for hyperspectral target detection. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 28(2), February 2006.
- [6] M. McCloud and L. Scharf. Interference estimation with applications to blind multiple-access communication over fading channels. *IEEE Transactions on Information Theory*, 46(3), May 2000.
- [7] C. McDiarmid. On method of bounded differences. *Surveys in Combinatorics*, 141:148–188, 1989.
- [8] M. Rangaswamy, F. Lin, and K. Gerlach. Robust adaptive signal processing methods for heterogeneous radar clutter scenarios. *Signal Processing*, 84:1653–1665, September 2004.
- [9] B. Recht. A simpler approach to matrix completion. 2009. Preprint available at <http://arxiv.org/abs/0910.0651>.
- [10] L. Scharf. *Statistical Signal Processing*. Addison-Wesley, Reading, MA, 1991.
- [11] L. Scharf and B. Friedlander. Matched subspace detectors. *IEEE Transactions on Signal Processing*, 42(8):2146–2157, August 1994.
- [12] D. Stein, S. Beaven, L. Hoff, E. Winter, A. Schaum, and A. Stocker. Anomaly detection from hyperspectral imagery. *IEEE Signal Processing Magazine*, January 2002.